

PEN TEST-D4D (MD-1)

Please read instructions carefully before use.

1)	Cap Dopper Plastic tube Collection paper Take all content out of the pouch.	6	Remove cap, pointing the tube away from your body. Point the tube down and apply few drops on the paper to cover the sampling area.	
2)	Take the (#633) test paper. Peel away the protective silicon cover. Hold it at the edge. Do not touch the exposed surface to avoid precontamination	7)	Check and compare results carefully with the color chart. The reaction for PCP and Methamphetamines may take up to 90 seconds to fully develop. Discard the test ingredients.	
3)	Wipe the suspected area, or touch the suspected substance with the test paper.	8)	Cannabis	
			PCP	
4)	Holding the tube upright, squeeze the plastic tube to break the inside ampoules. Do not squeeze tube again with broken glass inside.		Heroin/LSD/Opium/Morphine/ Methadone/Codeine	
(T)			Methamphetamine/Pethidine/Cathinone/ Ketamine/Ephedrine/Pseudoephedrine	
5)	Hold the cap with your finger and shake well for 5 seconds.		Amphetamine	
			No reaction	

Only colors similar to the color chart should be considered as positive results.

For each test a new test paper must be used. Use the test paper enclosed in each pouch.

Caution: keep out of reach of children. In case of eye or skin contact, flush well with water.

- 1. This pouch contains two tests. Once ampoule is broken, the test should be used within 1 hour.
- 2. Any positive color reactions may take 90 seconds or slightly longer to develop fully, depending on the quantity and purity of the substance being tested.
- 3. If there is no reaction to D4D (MD-1) or if the color reaction does not match the color chart, you can proceed to C&H (MD-2) which is the second Narcotic screening Test.

LII01708 V07